



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.


We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

same. The electrodes must be bright, as otherwise the phenomenon becomes very complicated. Furthermore the original polarization must often surpass a certain value if the residual polarization is to be contrary in sign; and there are other differences in detail for which there is no room here. Thus the rotational effect may proceed gradually to a maximum; an electromotive force zero may imply a very large residual polarization appearing on motion. The charging of moving electrodes is an interesting case; etc.

To elucidate this phenomenon, it suffices here to assume the occurrence of paired double layers $a\ a'$ and $b'\ b$, one double layer at each electrode. One element, a' , b' , of each double layer is localized in the liquid and the other element a , b in the solid electrode, both of the double layers having the same direction; *i. e.*, being two condensers in series. Hence there are two interpenetrating electrostatic fields, one



$b'\ a'$ localized in the liquid and the other $a\ b$ in the electrodes. These fields are in a contrary direction and the liquid field must be very much stronger to correspond with the initial counter polarization. On rotating the electrodes, the field localized in the liquid $b'\ a'$ is set free and its ions dissipated. The field localized in the solid, $a\ b$, however, remains and this constitutes the residual polarization in the direction of the charging current. Both fields decay in the lapse of time in the usual way.

When rotation ceases a liquid field is re-established, but usually, though not always, to a smaller degree. Eventually a probably discharges a' and b , b' , one of the fields passing through zero first, so that the effect of rotation finally vanishes. I have met both with marked polarization which on rotation vanished, as well as with an apparent absence of polarization which on rotation became very marked.

To obtain moving electrodes as free from polarization disturbances at the contact with

a liquid, it is therefore prudent to capture both fields; *i. e.*, to leave the electrodes entirely without interferences. This may be done by surrounding each with a porous cup, closed and completely filled with an electrolyte, the terminals passing out through an insulating tube. The electrodes should moreover be fixed rigidly to the cup. Again since zinc electrodes soon tarnish in brine but remain bright in concentrated zinc sulphate solution, the latter is a preferable electrolyte and the cups may be submerged in brine or any other solution.

I therefore constructed two cup electrodes of the kind in question and placed them in the rotational apparatus as before. The original potential difference of the zincs was about .4 millivolt. After keeping the circuit closed over night this fell off to below .05 millivolt, and could be eliminated by exchanging the cup electrodes. Rotation of the apparatus, *i. e.*, an external current in the brine surrounding the cups, produced no appreciable effect. The electrodes were then charged with a current of .2 am. for 30 sec. The polarization remaining was now much less than above, throughout, beginning with about 5 millivolts which fell to .5 m.v. in 5 minutes and to .05 in a few hours. Rotation was ineffective through all stages of the decay. No doubt the simple electrode in which both the original and the residual polarizations have vanished would often suffice, but with greater uncertainty, because such electrodes can not be exchanged without danger as to modifying their value.

CARL BARUS

BROWN UNIVERSITY,
PROVIDENCE, R. I.

SCIENCE

A Weekly Journal devoted to the Advancement of
Science, publishing the official notices and pro-
ceedings of the American Association for
the Advancement of Science

Published every Friday by

THE SCIENCE PRESS

LANCASTER, PA.

GARRISON, N. Y.

NEW YORK, N. Y.

Entered in the post-office at Lancaster, Pa., as second class matter